

# REDUCED SEDENTARY TIME INTERVENTION FOR BREAST CANCER SURVIVORS: OBJECTIVELY-MEASURED OUTCOMES FOR ACTIVITY AND METABOLISM

Rachel K. Walker<sup>1</sup>, Richard Viskochil<sup>2</sup>, Ruth Barham<sup>3</sup>, Jeanne Gagliarucci<sup>3</sup>, Ann Marie Moraitis<sup>1</sup> and Grace Makari-Judson<sup>3,4</sup>  
<sup>1</sup>Univ. of Massachusetts-Amherst, <sup>2</sup>Univ. of Utah, <sup>3</sup>Baystate Medical Center, <sup>4</sup>Univ. of Massachusetts Medical School



## BACKGROUND

Physical activity (PA) promotion and sedentary behavior reduction among cancer survivors is a national priority<sup>1</sup> and the number of PA-based behavioral interventions has expanded considerably in recent years.<sup>2</sup>

There have been relatively few trials focused on reduction of sedentary time among cancer survivors due in part to past limitations related to precise quantitative measurement of sedentary behaviors.<sup>3</sup>

Many PA interventions rely on clinic-based coaching which is both time-intensive and unrealistic for many clinics.<sup>4</sup>

The purpose of this study was to investigate the feasibility and effects of a home based 6 week reduced sedentary time intervention (RSTI) in breast cancer survivors who had completed primary treatment.

ClinicalTrials.gov NCT02969291

## METHODS

- Phase 1 proof-of-concept/feasibility trial
- One Group Pre/Post-test Design

### ELIGIBILITY CRITERIA

Designed to select a sample of breast cancer survivors most likely to benefit from a home-based intervention designed to reduce sedentary behavior.


#### Inclusion Criteria:

- Stage I-III breast cancer survivors age 20-80 who have completed primary treatment greater than 6 months but less than 5 years. Patients may be on adjuvant hormonal therapy.
- BMI >25
- Less than 150 min/week moderate to vigorous exercise
- No gain or loss of >10% body weight over prior 6 months

#### Exclusion Criteria:

- Known diabetes
- Known coronary artery disease
- Pregnancy

Figure 1. Study Flow (PA=Physical Activity, SB=Sedentary Behavior)




## INTERVENTION

Figure 2. ActivPAL Accelerometer



Figure 3. Personalized Feedback & Suggestions for Environmental Modifications to Reduce Sedentarism (selected pages)



## Metabolic parameters pre- and post intervention

Table 3. Metabolic Outcomes

n=13	Pre-intervention	Post-intervention	(%) change	P-value
<b>Hormones and metabolites</b>				
Fasting glucose (mg/dl)	126.1 ± 44.1	127.4 ± 34.2	+1.0	0.87
1h glucose (mg/dl)	180.2 ± 95.8	188.5 ± 82.4	+4.6	0.51
Fasting insulin (uU/ml)	15.3 ± 6.2	17.5 ± 9.0	+14.4	0.25
1h insulin (uU/ml)	121.6 ± 61.8	123.1 ± 58.3	+ 1.2	0.84
Leptin (ng/ml)	55.5 ± 31.6	44.6 ± 26.0	-19.6	0.01*
C-reactive protein (ng/ml)	237.1 ± 257.4	225.3 ± 259.5	- 5.0	0.68
Triglyceride (mg/dl)	107.2 ± 58.8	112.8 ± 52.8	+ 5.2	0.49
Total Cholesterol (mg/dl)	189.9 ± 22.2	195.6 ± 18.4	+ 3.0	0.32
HDL Cholesterol (mg/dl)	52.9 ± 14.6	58.5 ± 20.6	+ 10.6	0.07
LDL Cholesterol (mg/dl)	114.9 ± 26.6	114.6 ± 26.9	- 0.2	0.95
<b>Metrics of glycemic control</b>				
HOMA-IR	2.2 ± 1.1	2.4 ± 1.3	+ 9.0	0.34
HOMA-%B	86.9 ± 22.7	89.1 ± 24.1	+ 2.1	0.75
IGI (0-60)	0.40 ± 0.97	0.57 ± 0.73	+ 42.5	0.34
DI (HOMA-IR x IGI)	1.26 ± 2.69	1.56 ± 2.19	+23.8	0.51

Mean ± SD. HDL- High density lipoprotein, LDL- Low density lipoprotein, IGI- Insulinogenic index; DI- Disposition

Index. \*p<0.05

## RESULTS

Table 1. Participant Demographics (Total N=16)

Participants		
No. Completed study		13
Race	Black	3
	White	13
Median Age (range)		61.3 (49-73)
Median Time since diagnosis (range)		34.4 months (11-55)
Prior Therapy	Radiation	13
	Chemotherapy	11
	Hormonal	12

Table 2. Sedentary & Activity Outcomes

Variable Name	Average (SD)	T-Test (* = p < .05)
Total daily steps	Baseline: 6190.40 (2086.25)	0.76
	Post Intervention: 6326.30 (2788.41)	
Total energy expenditure/day (MET/hr.)	Baseline: 33.05 (0.92)	0.74
	Post: 33.11 (1.16)	
% of hourly time spent in uninterrupted sedentary behavior (8am-8pm)	Baseline: 20% (13%)	0.63
	Post Intervention: 22% (11%)	

## CONCLUSION

Results indicate that similar home-based RSTIs are safe, acceptable to survivors, and feasible to implement by cancer center staff.

Further research with larger samples and possible monitoring of interruptions in total sedentary time may be needed to establish efficacy and effect sizes for the intervention.

**A larger dose or addition of behavior-activating components (use of daily activity trackers, text messages, or coaching) may be necessary to realize clinically- meaningful changes in sedentarism, daily activity, metabolism and behavior change. These preliminary results suggest provision of educational material/one-time feedback is likely insufficient to meet PA guidelines & reduce sedentary time.**

➤ Acknowledgement of funding support from the Rays of Hope Foundation & the Susan G. Komen Foundation for Breast Cancer Research (CCR16376422)

➤ Acknowledgement of UMASS-Amherst nurses L.Carvalho, E.Lamoureux, and K. Bobianski for their contributions

- Schmitz KH et al. *American College of Sports Medicine roundtable on exercise guidelines for cancer survivors.* Med Sci. Sports Exerc. 42:1409-28, 2010.
- Phillips SM et al. *Physical activity and sedentary behavior in breast cancer survivors: new insights into activity patterns and potential intervention targets.* Gynecol. Oncol. 2015.
- Lynch BM et al. *Objectively measured physical activity and sedentary time of breast cancer survivors and associations with adiposity: findings from NHANES.* Cancer Causes Control 21:2:283-8, 2010.
- Courneya KS et al. *Subgroup effects in a randomized trial of different types and doses of exercise during breast cancer chemotherapy.* Br. J. Cancer 111:9: 1718-25, 2014.